

Deaths Due To Certain Conditions Originating In The Perinatal Period Among San Diego County Residents, 2003-2007

By Region, Race/Ethnicity, and Age Group

| | 2003 | | 2004 | | 2005 | | 2006 | | 2007 | |
|-------------------------------------|--------|--------------------------------|--------|--------------------------------|--------|--------------------------------|--------|--------------------------------|--------|--------------------------------|
| | Deaths | Adjusted Rate ^{a,b} | Deaths | Adjusted Rate ^{a,b} | Deaths | Adjusted Rate ^{a,b} | Deaths | Adjusted Rate ^{a,b} | Deaths | Adjusted Rate ^{a,b} |
| San Diego County^c | 113 | 3.7 | 128 | 4.0 | 129 | 4.0 | 114 | 3.4 | 126 | 3.8 |
| Region | | | | | | | | | | |
| N. Coastal | 11 | * | 22 | 3.7 | 20 | 3.3 | 26 | 4.3 | 17 | * |
| N. Central | 16 | * | 19 | * | 12 | * | 20 | 3.7 | 18 | * |
| Central | 20 | 3.9 | 23 | 4.2 | 21 | 3.8 | 22 | 4.0 | 23 | 4.1 |
| South | 23 | 4.8 | 21 | 3.9 | 29 | 5.2 | 17 | * | 32 | 5.3 |
| East | 14 | * | 20 | 4.4 | 12 | * | 10 | * | 13 | * |
| N. Inland | 27 | 5.3 | 21 | 3.9 | 33 | 6.0 | 17 | * | 23 | 4.0 |
| Gender | | | | | | | | | | |
| Male | 68 | 4.5 | 70 | 4.4 | 73 | 4.5 | 65 | 3.9 | 66 | 3.9 |
| Female | 45 | 3.0 | 58 | 3.6 | 56 | 3.5 | 49 | 3.0 | 60 | 3.6 |
| Race/Ethnicity^d | | | | | | | | | | |
| Hispanic | 50 | 4.3 | 48 | 3.7 | 63 | 4.8 | 45 | 3.2 | 54 | 3.8 |
| White | 30 | 2.4 | 43 | 3.4 | 32 | 2.5 | 39 | 3.1 | 34 | 2.7 |
| Black | 19 | * | 12 | * | 15 | * | 12 | * | 14 | * |
| Asian | 6 | * | 8 | * | 10 | * | 9 | * | 6 | * |
| Other | 6 | * | 12 | * | 9 | * | 9 | * | 18 | * |
| Age Groups | Deaths | Age-specific rate ^a | Deaths | Age-specific rate ^a | Deaths | Age-specific rate ^a | Deaths | Age-specific rate ^a | Deaths | Age-specific rate ^a |
| 0-4 | 112 | 53.5 | 127 | 57.7 | 128 | 57.1 | 114 | 49.7 | 125 | 53.8 |
| 5-14 | 0 | * | 0 | * | 0 | * | 0 | * | 1 | * |
| 15-24 | 1 | * | 1 | * | 1 | * | 0 | * | 0 | * |
| 25-34 | 0 | * | 0 | * | 0 | * | 0 | * | 0 | * |
| 35-44 | 0 | * | 0 | * | 0 | * | 0 | * | 0 | * |
| 45-54 | 0 | * | 0 | * | 0 | * | 0 | * | 0 | * |
| 55-64 | 0 | * | 0 | * | 0 | * | 0 | * | 0 | * |
| 65-74 | 0 | * | 0 | * | 0 | * | 0 | * | 0 | * |
| 75-84 | 0 | * | 0 | * | 0 | * | 0 | * | 0 | * |
| 85+ | 0 | * | 0 | * | 0 | * | 0 | * | 0 | * |

Deaths due to certain conditions originating in the perinatal period include ICD-10 codes P00-P96 as the underlying cause of death

* Rate not calculated for fewer than 20 deaths in this category.

^a All rates are per 100,000 population. Rates may vary slightly from previously published data due to updated population figures.

^b Adjusted rates are adjusted to 2000 U.S. Standard Population.

^c Deaths with unknown age, region or race/ethnicity are reflected in the county total, but not in the individual categories. Category sums may not add up to county total.

^d Race/ethnicity data may be different from previously published data due to modifications in the collecting, reporting and coding of racial and ethnic status.

Source: California Department of Public Health, Center for Health Statistics, Death Statistical Master Files;

SANDAG January 1 population estimates (2002-03 estimates rel. 2006, 2004-06 estimates rel. 2007, 2005-07 estimates rel. 2008)

Prepared by Community Epidemiology, Health & Human Services Agency, County of San Diego, 03/19/2009